

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

813 August 28, 1896

Reports of States and yearly and monthly reports of cities.

ALABAMA—Mobile.—Month of July, 1896. Estimated population, 31,076. Total deaths, 78, including phthisis pulmonalis, 11; enteric fever, 5, and diphtheria, 1.

California.—Month of July, 1896. Reports to the State board of health from 45 cities, towns, and villages, having an aggregate population of 653,014, show a mortality of 911 deaths, including phthisis pulmonalis, 131; enteric fever, 14; scarlet fever, 1; diphtheria, 4; eroup, 4, and whooping cough, 5.

MASSACHUSETTS—North Attleboro.—Month of July, 1896. Estimated population, 7,000. Total deaths, 11, including phthisis pulmonalis, 2, and diphtheria, 1.

MICHIGAN.—Week ended August 5, 1896. Reports to the State board of health, Lansing, from 52 observers, indicate that influenza, cholera infantum, consumption, and enteric fever increased, and that tonsillitis and inflammation of bowels decreased in area of prevalence. Phthisis pulmonalis was reported present during the week at 217 places, enteric fever at 55, scarlet fever at 20, diphtheria at 20, measles at 18, and whooping cough at 15 places.

MISSOURI—Kansas City.—Month of July, 1896. Estimated population, 165,000. Total deaths, 182, including phthisis pulmonalis, 6; enteric fever, 4; diphtheria, 4; measles, 2; croup, 1, and whooping cough, 7.

OHIO—Cleveland.—Month of July, 1896. Estimated population, 330,279. Total deaths, 617, including phthisis pulmonalis, 29; enteric fever, 13; diphtheria, 2, and measles, 4.

RHODE ISLAND—Newport.—Month of July, 1896. Estimated population, 21,500. Total deaths, 32, including 4 from phthisis pulmonalis and 1 from measles.

MORTALITY TABLE, CITIES OF THE UNITED STATES.

Cities.	Week ended.	Population, U. S. Census of 1890.	Total deaths from all causes.	Deaths from—										
				Phthisis pul- monalis.	Yellow fever.	Smallpox.	Varioloid.	Cholera.	Typhus fever.	Enteric fever.	Scarlet fever.	Diphtheria.	Measles.	Whooping cough.
Allegheny, Pa Do		105, 287 105, 287	31 42	2						3		1 2		i
Do		105, 287	36	2						4		2		
Do		105, 287	41	2						4		1		
Do		105, 287	44	3						2				
Ashtabula, Ohio	do	8,338	0	·										
Baltimore, Md	do	434, 439	182	19						3		5		3
Binghamton, N. Y		35, 005	9	1										
Boston, Mass		448, 477	215	30						3		10	1	1
Braddock, Pa		8, 561	1		•••••			•••••					•••••	
_ Do		8,561	8		••••					1				
Brockton, Mass		27, 294	35	2	•••••					1		•••••		
Do		27, 294	27	2										5
Brooklyn, N. Y	Aug. 22	806, 343	530	42	••••••			· · · · · ·		7	1 1	10	3	9